

## COLONOSCOPY Prep Instructions: Prepopik Prep

Location: Berkeley Endoscopy Center at 1072 Wildewood Centre Drive, Columbia, SC

Date: \_\_\_\_\_ Arrival time: \_\_\_\_\_ am /pm

If you need to reschedule, cancel or have any questions, please contact 803-788-1100 option 6.

**Please call 803-719-5242 to let us know you have arrived for your procedure.**

### **Key Instructions:**

Your doctor has scheduled you for a colonoscopy. To have a successful colonoscopy your colon must be clear of any stool. Follow all of the instructions in this handout **EXACTLY** as they are written. If you do not follow the directions, your procedure may be cancelled.

- **DO NOT** eat any solid foods after the light breakfast.
- Buy your bowel preparation at least **5 days** before your colonoscopy.
- **Do NOT** mix the solution until the day before your colonoscopy.
  
- A responsible family member or friend **MUST** drive you home from the procedure. You are **NOT ALLOWED** to drive, take a taxi, bus or car service, or leave the Endoscopy Center **ALONE**. If you do not have a responsible driver (family member or friend) with you to take you home, your procedure cannot be done with sedation and may be canceled. You **MUST** remain in the endoscopy center until you recover from sedation. **The entire process from arrival to discharge can take 3-4 hours. Please plan accordingly, as delays can often occur.**
- Wear comfortable, loose fitting clothing.
- Please bring with you any inhalers.
- Please do not bring any jewelry or other valuable items to the endoscopy center for your procedure.
- Wear comfortable, loose fitting clothing.
- Please bring with you any inhalers.
- Please do not bring any jewelry or other valuable items to the endoscopy center for your procedure.

### **Medications:**

Some of the medicines you take may need to be stopped or adjusted temporarily before colonoscopy

#### **3-5 Days before your Colonoscopy:**

- Stop Aspirin **such as** Ibuprofen.
  
- Stop Iron Supplements-**such as** Tandum Plus, Bifera, Ferrous Sulfate, Folic acid, Integra Plus
  
- Stop Arthritis medication such as Celebrex, Mobic, Ultram, Voltaren, Plaquenil, Naproxen). Do **NOT** take over the counter Aleve or Motrin.

## COLONOSCOPY Prep Instructions: Prepopik Prep

Location: Berkeley Endoscopy Center at 1072 Wildewood Centre Drive, Columbia, SC

Date: \_\_\_\_\_ Arrival time: \_\_\_\_\_ am /pm

You must speak to the prescribing physician on when you can stop taking these medications. Please do not stop taking without the consent of the physician.

- Blood thinners- **such as** Coumadin\* (warfarin), Plavix (clopidogrel), Ticlid (ticlopidine hydrochloride), Agrylin (anagrelide), Xarelto (Rivaroxaban), Pradaxa (Dabigatran), Eliquis (Apixaban), and Effient (Prasugrel).

### **Insulin and diabetes medications**

- Please call the doctor that monitors your glucose levels. Your medications may need to be adjusted due to the diet restrictions required with this bowel preparation. The following is some information to guide you:
- If you take a pill to lower your sugar, do not take it on the day of your procedure.
- If you are taking regular insulin, do not take it on the day of your procedure.
- If you taking long-acting insulin (NPH), take half of your prescribed dose on the morning of the procedure.
- If you are taking any other insulin preparation, such as Lantus, Humalog or 70/30 insulin, contact your prescribing physician for instructions.
- Please **take ALL** other prescribed medicines with a sip of water on **the day of your colonoscopy**.

### **You will need to pick up the following prescription from your pharmacy):**

- One (1) Prepopik Perp Kit
- One (1) Zofran ODT tablet –for nausea

### **You will need to buy the following over-the-counter (no prescription is needed).**

- Docolax (bisacodyl) laxative (not suppository or a stool-softener, you will need 4 tablets for the prep). You may choose a generic brand. **Please find this in the laxative aisle of the store.**

### **THREE (3) DAYS BEFORE YOUR COLONOSCOPY**

Avoid high-fiber foods-such as popcorn, beans, seeds (flax, sunflower and quinoa), multigrain bread, nuts, salad/vegetables, fresh and dried fruit. Avoid fiber supplements –such as Metamucil, Citrucel.

### **ONE (1) DAY BEFORE YOUR PROCEDURE**

- DO NOT eat any solid foods. The **clear liquids** you can drink include:
- Water, apple or white grape juice; fat free broth; coffee or tea (without milk or creamer); clear carbonated beverages such as ginger ale or lemon-lime soda; Gatorade or other sports drinks (not red or orange); Kool-Aid or other flavored drinks (**not red, purple or**

COLONOSCOPY Prep Instructions: Prepopik Prep

Location: Berkeley Endoscopy Center at 1072 Wildewood Centre Drive, Columbia, SC

Date: \_\_\_\_\_ Arrival time: \_\_\_\_\_ am /pm

**orange colors**). You may eat plain jello or gelatins (not red or orange) or popsicles (**not red, purple or orange colors**).

- **IMPORTANT TO KEEP HYDRATED**
- **Do NOT** drink alcohol on the day before or the day of the procedure or smoke cannabis (marijuana), as this can cause interaction with sedation.

**When to Drink Your Bowel Prep**


The day before your procedure:

<b>PACKET</b>  <b>1</b>	<b>STEP ONE</b>	<b>STEP TWO</b>
The afternoon of the day before your procedure	_____ <b>Fill</b> the dosing cup provided with <b>cold</b> water to the lower (5-ounce) line.	<b>Follow</b> with FIVE (5) 8-ounce drinks of clear liquid, taken at your own pace within the next 5 hours.
Date:  ____ / ____ / ____	_____ <b>Pour</b> in the contents of ONE (1) packet and <b>stir for 2-3 minutes until dissolved</b>	8 ounces 
Start Time: _____ PM  <b>(Between 4-6 PM)</b>	_____ <b>Drink</b> the entire contents  _____ <b>Take One Zofran tablet</b>	<b>See approved clear liquids above.</b>
<b>IMPORTANT</b> Take all your medications except for those on page 1 and 2.  Anal irritation or flare of hemorrhoid inflammation may occur. If this happens, treat it with over the counter remedies, such as hydro-cortisone cream, baby wipes, Vaseline, or TUCKS pads.	5 ounces - ] 	

COLONOSCOPY Prep Instructions: Prepopik Prep

Location: Berkeley Endoscopy Center at 1072 Wildewood Centre Drive, Columbia, SC

Date: \_\_\_\_\_ Arrival time: \_\_\_\_\_ am /pm

<b>PACKET</b>		
<b>2</b>		
	STEP ONE	STEP TWO
The late evening of the day before your colonoscopy:	_____ <b>Fill</b> the dosing cup provided with <b>cold</b> water to the lower (5-ounce) line.	<b>Follow</b> with THREE (3) 8-ounce drinks of clear liquid, taken at your own pace within the next 5 hours, <b>BEFORE</b> the colonoscopy.
Date: _____ / _____ / _____	_____ <b>Pour</b> in the contents of ONE (1) packet and <b>stir for 2-3 minutes until dissolved</b>	8 ounces ] 
Start Time: _____ PM ( Between 10PM -12AM )	_____ <b>Drink</b> the entire contents	<b>NOTE: You may continue to drink clear liquids until 2 hours before your colonoscopy.</b>
<b>REMINDER</b> Drink Clear liquids to stay hydrated up until 2 hours before colonoscopy.  <b>Take all your medications except for those on Page 1 and 2.</b>  <b>See approved clear liquids above.</b>	5 ounces -] 	